

Student's name

Student's number

Teacher's name



PLC PRESBYTERIAN
LADIES' COLLEGE
SYDNEY
1888

2014
TRIAL
HIGHER SCHOOL CERTIFICATE
EXAMINATION

Mathematics Extension 1

General Instructions

- Reading time - 5 minutes
- Working time - 2 hours
- Write using blue or black pen
Black is preferred
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- All necessary working should be shown in every question

Total Marks – 70

Section I: Pages 3-6

10 marks

- Attempt questions 1-10, using the answer sheet on page 13.
- Allow about 15 minutes for this section

Section II: Pages 7-10

60 marks

- Attempt questions 11-14, using the booklets provided.
- Allow about 1 hours 45 minutes for this section

Question	1-10	11	12	13	14	Total	%
Marks	/10	/15	/15	/15	/15	/70	

Blank Page

Section I


10 marks

Attempt Questions 1–10

Allow about 15 minutes for this section

Use the multiple-choice answer sheet for Questions 1–10

1.



What is the value of x ?

- (A) 1
- (B) $1\frac{1}{6}$
- (C) $2\frac{1}{6}$
- (D) $8\frac{1}{6}$

2.

What is the solution of $\frac{5}{1-x} < 3$?

- (A) $x < -\frac{2}{3}, x > 1$
- (B) $x < -\frac{2}{3}$
- (C) $x > 1$
- (D) $-\frac{2}{3} < x < 1$

3. What is the value of

$$\lim_{x \rightarrow 0} \left(\frac{2x}{\sin 5x} \right)?$$

(A) $\frac{2}{\sin 5}$

(B) $\frac{2}{5}$

(C) $\frac{5}{2}$

(D) $\sin 3x$

4. What are the co-ordinates of the point which divides the interval joining $A(3, -2)$ and $B(-5, 4)$ **externally** in the ratio of 5:3?

(A) $\left(0, \frac{1}{4} \right)$

(B) $(15, -11)$

(C) $(-17, 13)$

(D) $\left(-2, \frac{3}{2} \right)$

5. The inverse of the function $f(x) = e^{2x-1}$ is?


(A) $f^{-1}(x) = -e^{2x-1}$

(B) $f^{-1}(x) = \frac{e^{x+1}}{2}$

(C) $f^{-1}(x) = -\log_e(2x+1)$

(D) $f^{-1}(x) = \log_e \sqrt{x} + \frac{1}{2}$

6. The graph below shows a polynomial function, $y = P(x)$.



Which of the following could be the equation of $P(x)$?

- (A) $P(x) = (x+1)(x+2)(x+3)(x-1)$
 (B) $P(x) = -(x+1)(x+2)(x+3)(x-1)$
 (C) $P(x) = (x+1)(x-1)(x-2)(x-3)$
 (D) $P(x) = -(x+1)(x-1)(x-2)(x-3)$
7. The co-efficient of x^2 in the expansion $(2x-3)^5$ is?

- (A) -1080
 (B) -540
 (C) 540
 (D) 1080

8. Using $u = \cos x$,

$\int_0^{\frac{\pi}{3}} \sin^3 x \cos^4 x dx$ can be expressed in terms of u as

- (A) $\int_0^{\frac{\pi}{3}} u^6 - u^4 du$
 (B) $\int_0^1 u^6 - u^4 du$
 (C) $\int_{\frac{1}{2}}^1 u^4 - u^6 du$
 (D) $\int_0^{\frac{\sqrt{3}}{2}} u^4 - u^6 du$

9. A particle is moving along the x -axis, initially moving to the left from the origin. Its velocity and acceleration are given by


$$v^2 = 2 \log_e (3 + \cos x) \text{ and}$$

$$\ddot{x} = \frac{-\sin x}{3 + \cos x}.$$

Which of the following describes the subsequent motion?

- (A) Moves only to the left, alternately speeding up and slowing down, without becoming stationary.
- (B) Moves only to the left, alternately slowing to a stop and speeding up.
- (C) Slowing to a stop, then heading to the right forever.
- (D) Oscillates between two points.

10. Which of the following equations is shown in the sketch below?



- (A) $y = \cos^{-1}(\sin x)$
- (B) $y = \sin^{-1}(\cos x)$
- (C) $y = \sin^{-1}(x) + \sin(x)$
- (D) $y = \cos^{-1}(x) + \cos(x)$

Section II

60 marks

Attempt Questions 11–14

Allow about 1 hour and 45 minutes for this section.

Answer each question in a SEPARATE writing booklet. Extra writing booklets are available.

In Questions 11–14, your responses should include relevant mathematical reasoning and/or calculations.


Question 11 (15 marks) Use a SEPARATE writing booklet.

- a) Find the acute angle between the lines $x - 2y + 3 = 0$ and $y = 3x - 1$ at their point of intersection. **2**
- b) Find $\int \frac{1}{25 + 9x^2} dx$. **2**
- c) Find $\frac{d}{dx} \sin^{-1}(2x^3)$ **2**
- d) The polynomial $P(x) = x^3 - 3x^2 + kx + 12$ has 3 roots. It is known that two of the roots are of equal magnitude but opposite in sign. What is the value of k ? **3**
- e) Explain why Newton's method does not work for the root of the equation $x^3 - 3x + 6 = 0$ if the initial approximation is chosen to be $x = 1$. Use mathematics to support your answer. **2**
- f) If $\cot^2 \theta - \cot \theta = 1$, where $0 < \theta < \frac{\pi}{2}$,
- (i) Show that $\cot \theta = \frac{1 + \sqrt{5}}{2}$. **1**
- (ii) Hence, show that the exact value of $\cot 2\theta = \frac{1}{2}$. **3**

End of Question 11

Question 12 (15 marks) Use a SEPARATE writing booklet.

- a) AT is a tangent and is parallel to BP . Prove that $\angle ABP = \angle ACB$. 3



- b) A roast duck is taken out of the oven once it is cooked. A thermometer records the temperature of the duck to be 75°C . The roast duck is then allowed to cool in a room with a constant temperature of 23°C .
- (i) Show that $T = 23 + Ae^{-kt}$ satisfies the differential equation 1

$$\frac{dT}{dt} = -k(T - 23)$$
 where
 T is the temperature of the duck in degrees Celsius, $^{\circ}\text{C}$,
 t is the time in minutes and
 k is a constant.
- (ii) Show that $A = 52$. 1
- (iii) Find the value of k (in exact form) if after 5 minutes the duck's temperature is 65°C . 2
- (iv) Bacteria start to develop rapidly in the duck after 8 minutes. What will be the duck's temperature when the bacteria start to develop? Answer to the nearest degree. 1
- c) Using the substitution $u = e^x$, find $\int \frac{e^x dx}{\sqrt{1 - e^{2x}}}$ 2
- d) (i) Find the domain and range of the function $f(x) = \sin^{-1}(2x)$. 1
(ii) Sketch the graph of the function $f(x) = \sin^{-1}(2x)$. 1
(iii) The region bounded by the graph $f(x) = \sin^{-1}(2x)$ and the x -axis between $x = 0$ and $x = \frac{1}{2}$ is rotated about the y -axis to form a solid. 3
Find the exact volume of the solid.

End of Question 12

Question 13 (15 marks) Use a SEPARATE writing booklet.

- a)** The speed v m/s of a particle moving in a straight line is given by $v^2 = 84 + 16x - 4x^2$ where the displacement of the particle relative to a fixed point is x cm.
- (i) Find an expression for the particle's acceleration in terms of x . **2**
- (ii) Hence show that the particle is moving in simple harmonic motion. **1**
- (iii) Find the period, amplitude and centre of motion. **2**
- b)** (i) The monic polynomial, $P(x)$, has a root at $x = 3$, a double root at $x = -1$ and is of degree 4. If the polynomial passes through the point $(1,0)$, find the equation of the polynomial $P(x)$. **2**
- (ii) The polynomial $Q(x)$ has equation $Q(x) = x^2 + 1$. **2**
Show that $\frac{P(x)}{Q(x)}$ has a remainder of $4x + 8$.
- c)** A balloon has the shape of a right circular cylinder of radius r and length twice the radius, with a hemisphere at each end of radius r . The balloon is being filled at the rate of $10\text{cm}^3 / \text{s}$. Find the rate of change of r when $r = 8$ centimetres **2**
- d)** The points $P(2ap, ap^2)$ and $Q(2aq, aq^2)$ are the ends of a focal chord on the parabola $x^2 = 4ay$.
- (i) Show that PQ has equation $(p + q)x - 2y - 2apq = 0$. **1**
- (ii) Show that $pq = -1$ if PQ is a focal chord. **1**
- (iii) Show that the equation of the tangent at P is $y = px - ap^2$. **1**
- (iv) Hence find the locus of the point of intersection of the tangents at the ends of the focal chord. **1**

End of Question 13

Question 14 (15 marks) Use a SEPARATE writing booklet.


- a) Prove by Mathematical Induction that **3**

$$\sum_{r=1}^n \log_e \left(\frac{r+1}{r} \right) = \log_e (n+1)$$
 for all positive integers, n .

- b) Find the general solutions for $2 \cos x = \sqrt{3} \cot x$. **3**

- c) An object is projected horizontally from the top edge of a vertical cliff 40 metres above sea level with a velocity of 40 m/s .

Take $g = 10 \text{ m/s}^2$.



- (i) Using the top edge of the cliff as the origin, prove that the parametric equations of the path of the object are: **2**

$$x = 40t \qquad y = -5t^2 + 40$$
- (ii) Calculate when and where the object hits the water. **1**
- (iii) Find the velocity and angle of the object the instant it hits the water. **2**

- d) By considering $(1-x)^n \left(1 + \frac{1}{x}\right)^n$, or otherwise, express **4**

$$\binom{n}{2} \binom{n}{0} - \binom{n}{3} \binom{n}{1} + \dots + (-1)^n \binom{n}{n} \binom{n}{n-2}$$
 in simplest form.

End of Paper

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE : $\ln x = \log_e x, \quad x > 0$

Blank Page

Mathematics Extension 1:

Multiple Choice Answer Sheet

Student Number _____

Completely fill the response oval representing the most correct answer.

1. **A** **B** **C** **D**
2. **A** **B** **C** **D**
3. **A** **B** **C** **D**
4. **A** **B** **C** **D**
5. **A** **B** **C** **D**
6. **A** **B** **C** **D**
7. **A** **B** **C** **D**
8. **A** **B** **C** **D**
9. **A** **B** **C** **D**
10. **A** **B** **C** **D**

Blank Page