



SYDNEY BOYS HIGH  
MOORE PARK, SURRY HILLS

**2012**  
**TRIAL HIGHER SCHOOL**  
**CERTIFICATE EXAMINATION**

# Mathematics

## General Instructions:

- Reading time – 5 minutes
- Working time – 3 hours
- Write using black or blue pen  
Black pen is preferred
- Board approved calculators may be used
- A table of standard integrals is provided on the back of the Multiple Choice answer sheet
- All necessary working should be shown in every question if full marks are to be awarded
- Marks may NOT be awarded for messy or badly arranged work
- Answer in simplest exact form unless otherwise stated

## Total marks – 100 Marks

**Section I** Pages 2–4  
**10 marks**

- Attempt Questions 1–10
- Answer on the Multiple Choice answer sheet provided.
- Allow about 15 minutes for this section

**Section II** Pages 5–10  
**90 marks**

- Attempt Questions 11–16
- Allow about 2 hours 45 minutes for this section
- For Questions 11–16, start a new answer booklet per question

**Examiner:** Mr R. Boros

This is an assessment task only and does not necessarily reflect the content or format of the Higher School Certificate.

**Section I**— 10 marks

Select the alternative A, B, C, or D that best answers the question.

Fill in the response oval on your multiple choice answer sheet.

Marks

1.  $L + 2m - (L - 2n) - [2m + L - (2n - L)]$  simplifies to 1
- (A)  $4n + 2L$
- (B)  $4n - 2L$
- (C)  $4m + 4n - 2L$
- (D)  $4m - 4n - 2L$
2. At 10% p.a. simple interest, how long will it take for a sum of money to double? 1
- (A) 7.3 years
- (B) 5 years
- (C) 7.27 years
- (D) 10 years
3. What is the value of  $k$  if the expression  $4x^2 - 6x + k$  is a perfect square? 1
- (A)  $\frac{4}{9}$
- (B)  $\frac{9}{4}$
- (C) 4
- (D) 9
4.  $\frac{x^2 + 4x}{x^3 - 9x} \div \frac{x^2 + 2x - 8}{x^2 + x - 6}$  simplifies to 1
- (A) 1
- (B)  $\frac{x}{x - 3}$
- (C)  $\frac{1}{x - 3}$
- (D)  $\frac{1}{x + 3}$

5. The solution to the equation  $2x^2 = 7x$  is  $x =$


1

- (A) 0 or  $-3^{1/2}$
- (B) 0 or  $3^{1/2}$
- (C)  $3^{1/2}$  only
- (D)  $3^{1/2}$  or  $-3^{1/2}$

6. If  $p$  and  $q$  are the roots of  $15x^2 + 75x - 3 = 0$  then  $p + q =$

1

- (A) 75
- (B) 5
- (C)  $-\frac{1}{5}$
- (D) -5



1

The shaded region in the diagram satisfies

- (A)  $x + 2 \geq 2y$  and  $x + 2y > 2$
- (B)  $x + 2 \geq 2y$  and  $x + 2y < 2$
- (C)  $x + 2 \leq 2y$  and  $x + 2y > 2$
- (D)  $x + 2 \leq 2y$  and  $x + 2y < 2$

8.  $\log_3 15 + \log_3 18 - \log_3 10 =$

1

- (A) 1
- (B) 2
- (C) 3
- (D) 0

9. Two cards are drawn in succession from a regular pack of 52 cards. What is the probability that both cards are diamonds or both cards are clubs? 1

(A)  $\frac{2}{17}$

(B)  $\frac{3}{5}$

(C)  $\frac{3}{17}$

(D)  $\frac{27}{52}$

10. If the 5th term and 18th term of an arithmetic series are 12 and 64 respectively, find the common difference. 1

(A) -5

(B) 4

(C) -4

(D) 5

**Section II**— 90 marks

Marks

**Question 11 (15 marks)** (use a separate answer booklet)

(a) Find the first derivative of

(i)  $y = (x^2 - 1)^3$ , 2


(ii)  $y = \frac{2x}{x - 1}$ , 2

(iii)  $f(x) = \ln(3 - x)$ . 1

(b) Evaluate  $\int_3^8 \sqrt{x + 1} dx$ . 3

(c) Find the equation of the normal to the curve  $y = \tan x$  at the point where  $x = \frac{\pi}{4}$  (answer in the general form of a line). 3

(d) The graph shows  $y'$  and  $y''$  for the function  $y = f(x)$ . 3



Sketch a graph of  $y = f(x)$ , clearly showing the  $x$  values of any turning points and points of inflexion.


(e) Find  $\int 3 \cos\left(\frac{x}{2}\right) dx$ . 1

**Question 12 (15 marks)** (use a separate answer booklet)

(a)  $A(-1, 8)$ ,  $B(4, -2)$ , and  $C(-3, -1)$  are three points on the number plane. The line  $\ell_1$  passes through the points  $A$  and  $B$ .

- (i) Draw a sketch showing  $A$ ,  $B$ ,  $C$ , and  $\ell_1$ . 1
- (ii) Find the exact distance  $AB$ . 1
- (iii) Show that  $\ell_1$  has the equation  $2x + y - 6 = 0$ . 1
- (iv) Find the perpendicular distance from the point  $C$  to the line  $2x + y - 6 = 0$ . 2
- (v) Calculate the area of the triangle  $ABC$ . 2
- (vi) Find the co-ordinates of the midpoint,  $M$ , of  $AC$ . 1
- (vii) Find the equation of the line,  $\ell_2$ , through  $M$  and parallel to  $AB$  (written in the general form of a line). 2

(b) Two separate ‘one man’ canoes start off from a jetty,  $P$ , on a very large lake. The first canoeist paddles on a bearing of  $040^\circ \text{T}$  for 12 nautical miles to a buoy  $Q$ . At the same time the second canoeist paddles a distance of 8 nautical miles on a bearing of  $100^\circ \text{T}$  to another buoy  $R$ .



- (i) Copy the sketch above and add all the relevant information. 1
- (ii) Calculate the distance (in nautical miles) between the canoeists correct to one decimal place. 2
- (iii) If the two canoeists conduct a quick search of  $\triangle PQR$  for any other canoeists, calculate the total area searched, giving your answer in square kilometres correct to the nearest 10 square kilometres. (Note: 1 nautical mile = 1852 metres.) 2

**Question 13 (15 marks)** (use a separate answer booklet)

(a) (i) For the curve  $y = 3 \sin 4x$  in the domain  $0 \leq x \leq \pi$ , state the  
 ( $\alpha$ ) period, 1


( $\beta$ ) amplitude. 1

(ii) Sketch the curve  $y = 3 \sin 4x$ ,  $0 \leq x \leq \pi$ , clearly showing where the curve cuts the  $x$ -axis. 2

(iii) Hence or otherwise, find the NUMBER of solutions to  $\sin 4x = \cos x$  where  $0 \leq x \leq \pi$ . 2

(b) Evaluate  $\int_0^1 2xe^{(3x^2-5)} dx$ , giving your answer to 3 significant figures. 3

(c) A right circular cylinder of radius  $r$  cm and height  $h$  cm has to be designed to fit inside a sphere of  $2\sqrt{3}$  cm radius so that both the bottom and the top touch the sphere (centre  $O$ ) completely on the circular rim.



(i) Using the diagram as a guide, show that  $r^2 = 12 - \frac{h^2}{4}$ . 1

(ii) If the volume of the cylinder is  $V$ , show that  $V = 12\pi h - \frac{1}{4}\pi h^3$ . 1

(iii) Hence find the dimensions of the cylinder to give maximum volume. 2

(d) Evaluate  $\int_0^5 \frac{x}{5 + 3x^2} dx$ , leaving your answer in exact form. 2


**Question 14 (15 marks)** (use a separate answer booklet)

- (a) Given that  $x^2 - 6x - 7 = 8y$ , find:
- (i) the co-ordinates of the vertex, 1
  - (ii) the co-ordinates of the focus, 1
  - (iii) the equation of the directrix. 1
- (b) A road grader removes  $V \text{ m}^3$  of soil in  $t$  minutes, where  $V = 25t - \frac{t^2}{50}$ . 2  
Find the rate at which the soil is being removed after five minutes.
- (c) A driver in a car is at a point  $A$ , from which branches out two roads. If he takes the road on the **LEFT** and journeys some distance, this road leads to a point  $B$  from which branches off three roads, one of which leads to his destination  $C$ . However if he takes the road on the **RIGHT**, and journeys along a certain distance, this road leads to a point  $D$ , from which branches off four roads, one of which leads to his destination  $C$ . 3  
Assuming he has no Sat. Navigation or prior knowledge of any of these facts, except that he wants to travel to destination  $C$ , find the probability that he **DOES NOT** reach  $C$  on his first try.
- (d) Evaluate  $\int_0^{\frac{\pi}{2}} \sin^2 x \, dx + \int_0^{\frac{\pi}{2}} \cos^2 x \, dx$ . 2
- (e) Let  $A(4, 0)$  and  $B(1, 0)$  be two fixed points and let  $P$  be the variable point  $(x, y)$ .
- (i) Write down expressions for the distances  $PA$  and  $PB$  in terms of  $x$  and  $y$ . 2
  - (ii) Find the locus of  $P$  whose distance from  $A$  is twice its distance from  $B$ . 3



**Question 15 (15 marks)** (use a separate answer booklet)

- (a) The diagram below shows an ancient window which consists of a rectangle  $ABCD$  with height  $2\sqrt{3}$  m and width 2 m surmounted by a minor segment of a circle which is stained glass. The centre of the circle is at  $O$ , the point of intersection of the diagonals of the rectangle.



- (i) Explain why  $\widehat{AOB} = 60^\circ$ . 2
- (ii) Find the area of the minor segment correct to 3 decimal places. 2
- (b) (i) Sketch the region beneath the curve  $y = e^x + 1$  which is above the  $x$ -axis and between the lines  $x = 0$  and  $x = 1$ . 1
- (ii) The region in (b)(i) is now rotated about the  $x$ -axis. Find the volume of the resulting solid of revolution. Leave your answer in exact form. 3
- (c) For the curve  $y = xe^{-x}$ ,
- (i) Prove that  $\frac{dy}{dx} = -e^{-x}(x - 1)$ . 1
- (ii) Find any stationary points and determine their nature. 2
- (iii) Prove that  $\frac{d^2y}{dx^2} = e^{-x}(x - 2)$ . 1
- (iv) Show that there is a point of inflexion on this curve and find the co-ordinates of this point. 2
- (v) Sketch the curve, showing the co-ordinates of the point of inflexion and any stationary points. 1

**Question 16 (15 marks)** (use a separate answer booklet)

- (a) The number of DVD copies sold at a store of *The London Olympics 2012 Opening Ceremony* has increased exponentially in accordance with the formula  $N = Ae^{kt}$  where  $t$  is the time in weeks after the Opening Ceremony. Initially 10 000 copies were sold and the number doubled after two weeks.

(i) Find the value of  $A$ . 1

(ii) Calculate the value of  $k$  correct to 3 decimal places. 1

(iii) At what rate was the number of copies increasing after four weeks? Answer correct to the nearest whole number. 2

- (b) Mr B—— borrows  $\$P$  to fund his new Nissan supercar. The term of the loan is 10 years with an interest rate of 6% p.a., monthly reducible. He repays the loan in equal monthly installments of  $\$750$ .

(i) Show that at the end of  $n$  months, the amount owing is given by  $A = P(1.005)^n - 150\,000(1.005)^n + 150\,000$ . 3


(ii) If at the end of 10 years the loan has been repaid, calculate the amount that he originally borrowed, correct to the nearest dollar. 2

- (c) A particle is moving along the  $x$ -axis. The distance of the particle,  $x$  metres from the origin  $O$  is given by the equation  $x = 6t + e^{-4t}$  where  $t$  is time in seconds.

(i) Write down an expression for velocity of the particle. 1

(ii) Explain why the particle will never come to rest. 2

- (d)  $A$ ,  $B$ , and  $C$  are the vertices of an isosceles triangle where  $AC = BC$  and right-angled at  $C$ .  $D$  is a point such that  $DB = AB$  and  $\widehat{D\hat{B}A}$  is acute.  $DC \parallel AB$  also. 3



Find, giving reasons, the size of  $\widehat{D\hat{B}C}$ .

**End of Paper**

**Blank Page**

**Blank Page**

**Section I**— 10 marks

Select the alternative A, B, C, or D that best answers the question.

Fill in the response oval on your multiple choice answer sheet.

Marks

1.  $L + 2m - (L - 2n) - [2m + L - (2n - L)]$  simplifies to

- (A)  $4n + 2L$   
(B)  $4n - 2L$   
(C)  $4m + 4n - 2L$   
(D)  $4m - 4n - 2L$

1

**Solution:** 
$$L + 2m - L + 2n - (2m + L - 2n + L) = 2m + 2n - (2m - 2n + 2L) = 4n - 2L$$

2. At 10% p.a. simple interest, how long will it take for a sum of money to double?

- (A) 7.3 years  
(B) 5 years  
(C) 7.27 years  
(D) 10 years

1

**Solution:** 
$$I = P \times \frac{10}{100} \times n = P \text{ (i.e. } 2P = P + I)$$
$$\text{so } P = \frac{Pn}{10}$$
$$n = 10$$

3. What is the value of  $k$  if the expression  $4x^2 - 6x + k$  is a perfect square?

- (A)  $\frac{4}{9}$   
(B)  $\frac{9}{4}$   
(C) 4  
(D) 9

1

**Solution:** 
$$4\left(x^2 - \frac{6x}{4} + \frac{k}{4}\right) = 4\left(x^2 - \frac{3x}{2} + \frac{9}{16}\right)$$
$$\text{So } k = \frac{9}{4}$$

4.  $\frac{x^2 + 4x}{x^3 - 9x} \div \frac{x^2 + 2x - 8}{x^2 + x - 6}$  simplifies to

1

(A) 1

(B)  $\frac{x}{x-3}$

(C)  $\frac{1}{x-3}$

(D)  $\frac{1}{x+3}$

**Solution:** 
$$\frac{x(x+4)}{x(x^2-9)} \times \frac{(x+3)(x-2)}{(x+4)(x-2)} = \frac{(x+3)}{(x+3)(x-3)}$$

$$= \frac{1}{x-3}$$

5. The solution to the equation  $2x^2 = 7x$  is  $x =$

1

(A) 0 or  $-3^{1/2}$

(B) 0 or  $3^{1/2}$

(C)  $3^{1/2}$  only

(D)  $3^{1/2}$  or  $-3^{1/2}$

**Solution:**  $x(2x - 7) = 0$   
 $\therefore x = 0$  or  $3^{1/2}$

6. If  $p$  and  $q$  are the roots of  $15x^2 + 75x - 3 = 0$  then  $p + q =$

1

(A) 75


(B) 5

(C)  $-\frac{1}{5}$

(D) -5

**Solution:** 
$$p + q = -\frac{75}{15}$$

$$= -5$$



1

The shaded region in the diagram satisfies

- (A)  $x + 2 \geq 2y$  and  $x + 2y > 2$
- (B)  $x + 2 \geq 2y$  and  $x + 2y < 2$
- (C)  $x + 2 \leq 2y$  and  $x + 2y > 2$
- (D)  $x + 2 \leq 2y$  and  $x + 2y < 2$

**Solution:** Testing  $(2, 1)$  in  $x + 2 \geq 2y$  gives true,  
and testing  $(2, 1)$  in  $x + 2y > 2$  gives true,  
so the correct answer is (A).

8.  $\log_3 15 + \log_3 18 - \log_3 10 =$

1

- (A) 1
- (B) 2
- (C) 3
- (D) 0

**Solution:**

$$\begin{aligned} \log_3 \left( \frac{15 \times 18}{10} \right) &= \log_3 27 \\ &= \log_3 3^3 \\ &= 3 \end{aligned}$$

9. Two cards are drawn in succession from a regular pack of 52 cards. What is the probability that both cards are diamonds or both cards are clubs? 1

- (A)  $\frac{2}{17}$   
(B)  $\frac{3}{5}$   
(C)  $\frac{3}{17}$   
(D)  $\frac{27}{52}$

<p><b>Solution:</b> <math>P(\text{both diamonds}) = \frac{1}{4} \times \frac{12}{51} = \frac{1}{17}</math>, <math>\therefore P(\text{both diamonds or both clubs}) = \frac{1}{17} + \frac{1}{17} = \frac{2}{17}</math></p>
--

10. If the 5th term and 18th term of an arithmetic series are 12 and 64 respectively, find the common difference. 1

- (A) -5  
(B) 4  
(C) -4  
(D) 5

<p><b>Solution:</b> <math>18 - 5 = 13</math> <math>(64 - 12) \div 13 = 4</math></p>
---



**Section II**— 90 marks

Marks

**Question 11 (15 marks)** (use a separate answer booklet)

(a) Find the first derivative of

(i)  $y = (x^2 - 1)^3$ ,

2

**Solution:**  $3 \times 2x(x^2 - 1)^2 = 6x(x^2 - 1)^2$ .

(ii)  $y = \frac{2x}{x - 1}$ ,

2

**Solution:**  $\frac{(x - 1) \cdot 2 - 2x}{(x - 1)^2} = \frac{-2}{(x - 1)^2}$ .

(iii)  $f(x) = \ln(3 - x)$ .

1

**Solution:**  $\frac{-1}{3 - x}$ .

(b) Evaluate  $\int_3^8 \sqrt{x + 1} dx$ .

3

**Solution:**  $\int_3^8 (x + 1)^{1/2} \cdot dx = \left[ \frac{2(x + 1)^{3/2}}{3} \right]_3^8$ ,  
 $= 18 - \frac{16}{3}$ ,  
 $= \frac{38}{3}$  or  $12\frac{2}{3}$ .


(c) Find the equation of the normal to the curve  $y = \tan x$  at the point where  $x = \frac{\pi}{4}$  (answer in the general form of a line).

3


**Solution:** Point  $(\frac{\pi}{4}, 1)$ ,  
 $y' = \sec^2 x$ ,  $\therefore$  tangent slope = 2.  
Hence normal is:  $y - 1 = -\frac{1}{2}(x - \frac{\pi}{4})$ ,  
 $2y - 2 = -x + \frac{\pi}{4}$ ,  
 $x + 2y - 2 - \frac{\pi}{4} = 0$ .

- (d) The graph shows  $y'$  and  $y''$  for the function  $y = f(x)$ .

3



Sketch a graph of  $y = f(x)$ , clearly showing the  $x$  values of any turning points and points of inflexion.



- (e) Find  $\int 3 \cos\left(\frac{x}{2}\right) dx$ .

1

**Solution:**


$$\begin{aligned} \int 3 \cos\left(\frac{x}{2}\right) dx &= 2 \times 3 \sin\left(\frac{x}{2}\right) + c, \\ &= 6 \sin\left(\frac{x}{2}\right) + c, \end{aligned}$$

**Question 12 (15 marks)** (use a separate answer booklet)

- (a)  $A(-1, 8)$ ,  $B(4, -2)$ , and  $C(-3, -1)$  are three points on the number plane. The line  $\ell_1$  passes through the points  $A$  and  $B$ .

- (i) Draw a sketch showing  $A$ ,  $B$ ,  $C$ , and  $\ell_1$ .

1



- (ii) Find the exact distance  $AB$ .

1

**Solution:**

$$\begin{aligned}
 AB &= \sqrt{(4 + 1)^2 + (-2 - 8)^2}, \\
 &= \sqrt{25 + 100}, \\
 &= 5\sqrt{5}.
 \end{aligned}$$

- (iii) Show that  $\ell_1$  has the equation  $2x + y - 6 = 0$ .

1

**Solution:**

$$\begin{aligned}
 \text{Slope} &= \frac{8 + 2}{-1 - 4}, \\
 &= -2.
 \end{aligned}$$

Equation:  $y + 2 = -2(x - 4)$ ,

$$\begin{aligned}
 &= -2x + 8, \\
 \therefore 2x + y - 6 &= 0.
 \end{aligned}$$

- (iv) Find the perpendicular distance from the point  $C$  to the line  $2x + y - 6 = 0$ .

2

**Solution:**

$$\begin{aligned}
 \text{Perp. distance} &= \frac{|2(-3) + (-1) - 6|}{\sqrt{4 + 1}}, \\
 &= \frac{13}{\sqrt{5}} \text{ or } \frac{13\sqrt{5}}{5}.
 \end{aligned}$$

- (v) Calculate the area of the triangle  $ABC$ .

2

$$\begin{aligned} \text{Solution: Area} &= \frac{1}{2} \times 5\sqrt{5} \times \frac{14}{\sqrt{5}}, \\ &= \frac{65}{2}. \end{aligned}$$

- (vi) Find the co-ordinates of the midpoint,  $M$ , of  $AC$ .

1


$$\begin{aligned} \text{Solution: } M &= \left( \frac{-1-3}{2}, \frac{8-1}{2} \right), \\ &= (-2, 7/2). \end{aligned}$$

- (vii) Find the equation of the line,  $l_2$ , through  $M$  and parallel to  $AB$  (written in the general form of a line).

2

$$\begin{aligned} \text{Solution: } l_2 \text{ is } y - \frac{7}{2} &= -2(x + 2), \\ 2x + y + 1/2 &= 0. \end{aligned}$$


- (b) Two separate 'one man' canoes start off from a jetty,  $P$ , on a very large lake. The first canoeist paddles on a bearing of  $040^\circ$  T for 12 nautical miles to a buoy  $Q$ . At the same time the second canoeist paddles a distance of 8 nautical miles on a bearing of  $100^\circ$  T to another buoy  $R$ .



- (i) Copy the sketch above and add all the relevant information.

1

**Solution:**



- (ii) Calculate the distance (in nautical miles) between the canoeists correct to one decimal place. 2

**Solution:**  $p^2 = 8^2 + 12^2 - 2 \cdot 8 \cdot 12 \cdot \cos 60^\circ,$   
 $= 112.$   
 $\therefore$  Distance between is 10.6 n.m.

- (iii) If the two canoeists conduct a quick search of  $\triangle PQR$  for any other canoeists, calculate the total area searched, giving your answer in square kilometres correct to the nearest 10 square kilometres. (Note: 1 nautical mile = 1852 metres.) 2

**Solution:** Area =  $\frac{1}{2} \times 8 \times 12 \times \sin 60^\circ \times (1.852)^2,$   
 $\approx 140 \text{ km}^2$  (nearest 10  $\text{km}^2$ ).

**Question 13 (15 marks)** (use a separate answer booklet)

- (a) (i) For the curve  $y = 3 \sin 4x$  in the domain  $0 \leq x \leq \pi$ , state the  
 (α) period,

1

**Solution:**  $\frac{\pi}{2}$ .


- (β) amplitude.

1

**Solution:** 3.

- (ii) Sketch the curve  $y = 3 \sin 4x$ ,  $0 \leq x \leq \pi$ , clearly showing where the curve cuts the  $x$ -axis.

2



- (iii) Hence or otherwise, find the NUMBER of solutions to  $\sin 4x = \cos x$  where  $0 \leq x \leq \pi$ .

2


**Solution:** Multiply throughout by 3 gives  $3 \sin 4x = 3 \cos x$ , then graph  $y = 3 \cos x$  on the diagram above. It is clear that there are 5 solutions.

- (b) Evaluate  $\int_0^1 2xe^{(3x^2-5)} dx$ , giving your answer to 3 significant figures.

3

**Solution:** Note that  $\frac{d}{dx} (e^{(3x^2-5)}) = 6xe^{(3x^2-5)}$ .  
 So  $\frac{1}{3} \int_0^1 6xe^{(3x^2-5)} dx = \frac{1}{3} [e^{(3x^2-5)}]_0^1$ ,  
 $= \frac{1}{3} (e^{-2} - e^{-5})$ ,  
 $\approx 0.0429$  (3 sig. fig.)

- (c) A right circular cylinder of radius  $r$  cm and height  $h$  cm has to be designed to fit inside a sphere of  $2\sqrt{3}$  cm radius so that both the bottom and the top touch the sphere (centre  $O$ ) completely on the circular rim.



- (i) Using the diagram as a guide, show that  $r^2 = 12 - \frac{h^2}{4}$ . 1

**Solution:**  $\left(\frac{h}{2}\right)^2 + r^2 = (2\sqrt{3})^2$  (by Pythagoras's Thm.),

$$r^2 = 4 \times 3 - \frac{h^2}{4},$$

$$= 12 - \frac{h^2}{4}.$$

- (ii) If the volume of the cylinder is  $V$ , show that  $V = 12\pi h - \frac{1}{4}\pi h^3$ . 1

**Solution:**  $V = \pi r^2 h,$

$$= \pi h \left(12 - \frac{h^2}{4}\right),$$

$$= 12\pi h - \frac{\pi h^3}{4}.$$

(iii) Hence find the dimensions of the cylinder to give maximum volume.

2

**Solution:**

$$\begin{aligned}\frac{dV}{dh} &= 12\pi - \frac{3\pi h^2}{4}, \\ &= 0 \text{ when } 12\pi = \frac{3\pi h^2}{4}, \\ &\quad 48\pi = 3\pi h^2, \\ &\quad h^2 = 16, \\ &\quad h = 4 \text{ cm.}\end{aligned}$$
$$\begin{aligned}\frac{d^2V}{dh^2} &= -\frac{6\pi h}{4}, \\ &= -\frac{3\pi h}{2}, \\ &= -6\pi \text{ when } h = 4, \\ &< 0 \implies \text{maximum.}\end{aligned}$$

So  $r^2 = 12 - \frac{16}{4}$ ,

$$\begin{aligned}&= 8, \\ r &= 2\sqrt{2} \text{ cm.}\end{aligned}$$

(d) Evaluate  $\int_0^5 \frac{x}{5+3x^2} dx$ , leaving your answer in exact form.

2

**Solution:**

$$\begin{aligned}\frac{1}{6} \int_0^5 \frac{6x}{5+3x^2} dx &= \frac{1}{6} \left[ \ln(5+3x^2) \right]_0^5, \\ &= \frac{1}{6} \ln \frac{80}{5}, \\ &= \frac{1}{3} \ln 4.\end{aligned}$$



**Question 14 (15 marks)** (use a separate answer booklet)

(a) Given that  $x^2 - 6x - 7 = 8y$ , find:

(i) the co-ordinates of the vertex, 1

**Solution:**  $x^2 - 6x + 3^2 = 8y + 7 + 9,$   
 $(x - 3)^2 = 4 \times 2(y + 2).$   
 So the vertex is  $(3, -2).$

(ii) the co-ordinates of the focus, 1

**Solution:** From above, the focus is  $(3, 0).$

(iii) the equation of the directrix. 1

**Solution:** Also from above, the directrix is  $y = -4.$

(b) A road grader removes  $V \text{ m}^3$  of soil in  $t$  minutes, where  $V = 25t - \frac{t^2}{50}$ . 2  
 Find the rate at which the soil is being removed after five minutes.

**Solution:**  $\frac{dV}{dt} = 25 - \frac{2t}{50}.$   
 When  $t = 5,$   $\frac{dV}{dt} = 25 - \frac{5}{25},$   
 $= 24\frac{4}{5}.$   
 $\therefore$  The rate of removal is  $24\frac{4}{5} \text{ m}^3/\text{min}.$

(c) A driver in a car is at a point  $A$ , from which branches out two roads. If he takes the road on the LEFT and journeys some distance, this road leads to a point  $B$  from which branches off three roads, one of which leads to his destination  $C$ . However if he takes the road on the RIGHT, and journeys along a certain distance, this road leads to a point  $D$ , from which branches off four roads, one of which leads to his destination  $C$ . 3

Assuming he has no Sat. Navigation or prior knowledge of any of these facts, except that he wants to travel to destination  $C$ , find the probability that he DOES NOT reach  $C$  on his first try.

**Solution:**  $P(C \text{ via } B) = \frac{1}{2} \times \frac{1}{3},$   
 $= \frac{1}{6}.$   
 $P(C \text{ via } D) = \frac{1}{2} \times \frac{1}{4},$   
 $= \frac{1}{8}.$   
 $P(\tilde{C}) = 1 - \left(\frac{1}{6} + \frac{1}{8}\right),$   
 $= \frac{17}{24}.$

- (d) Evaluate  $\int_0^{\frac{\pi}{2}} \sin^2 x \, dx + \int_0^{\frac{\pi}{2}} \cos^2 x \, dx$ .

2

**Solution:** 
$$\begin{aligned} \int_0^{\frac{\pi}{2}} \sin^2 x \, dx + \int_0^{\frac{\pi}{2}} \cos^2 x \, dx &= \int_0^{\frac{\pi}{2}} 1 \, dx, \\ &= x \Big|_0^{\frac{\pi}{2}}, \\ &= \frac{\pi}{2}. \end{aligned}$$

- (e) Let  $A(4, 0)$  and  $B(1, 0)$  be two fixed points and let  $P$  be the variable point  $(x, y)$ .

- (i) Write down expressions for the distances  $PA$  and  $PB$  in terms of  $x$  and  $y$ .

2

**Solution:**  $PA = \sqrt{(x-4)^2 + y^2}, \quad PB = \sqrt{(x-1)^2 + y^2}$

- (ii) Find the locus of  $P$  whose distance from  $A$  is twice its distance from  $B$ .


3

**Solution:** 
$$\begin{aligned} PA &= 2PB, \\ PA^2 &= 4PB^2, \\ (x-4)^2 + y^2 &= 4\{(x-1)^2 + y^2\}, \\ x^2 - 8x + 16 + y^2 &= 4\{x^2 - 2x + 1 + y^2\}, \\ 3x^2 + 3y^2 &= 12, \\ x^2 + y^2 &= 2^2. \end{aligned}$$

So the locus is a circle with centre at the origin and radius 2.

**Question 15 (15 marks)** (use a separate answer booklet)

- (a) The diagram below shows an ancient window which consists of a rectangle  $ABCD$  with height  $2\sqrt{3}$  m and width 2 m surmounted by a minor segment of a circle which is stained glass. The centre of the circle is at  $O$ , the point of intersection of the diagonals of the rectangle.



- (i) Explain why  $\widehat{AOB} = 60^\circ$ .

2


**Solution:**  $DB = \sqrt{4 + 12}$ ,  
 $= 4$  m.  
 $O$  bisects  $DB$ ,  $\therefore OB = 2$  m.  
 Similarly  $OB = 2$  m and,  $AB = 2$  m,  
 $\triangle OAB$  is equilateral, so  $\widehat{AOB} = 60^\circ$ .

- (ii) Find the area of the minor segment correct to 3 decimal places.

2

**Solution:** Minor segment = sector  $AOB - \triangle AOB$ ,  
 $= \frac{1}{6} \times \pi \times 2^2 - \frac{1}{2} \times 2^2 \times \sin 60^\circ$ ,  
 $\approx 0.362 \text{ m}^2$  (3 d.p.)

- (b) (i) Sketch the region beneath the curve  $y = e^x + 1$  which is above the  $x$ -axis and between the lines  $x = 0$  and  $x = 1$ . 1



- (ii) The region in (b)(i) is now rotated about the  $x$ -axis. Find the volume of the resulting solid of revolution. Leave your answer in exact form. 3

**Solution:**

$$\begin{aligned} \text{Vol.} &= \pi \int_0^1 y^2 dx, \\ &= \pi \int_0^1 (e^{2x} + 2e^x + 1) dx, \\ &= \pi \left[ \frac{e^{2x}}{2} + 2e^x + x \right]_0^1, \\ &= \pi \left\{ \frac{e^2}{2} + 2e + 1 - \left( \frac{1}{2} + 2 + 0 \right) \right\}, \\ &= \pi \left( \frac{e^2}{2} + 2e - \frac{3}{2} \right). \end{aligned}$$

- (c) For the curve  $y = xe^{-x}$ ,

- (i) Prove that  $\frac{dy}{dx} = -e^{-x}(x - 1)$ . 1

**Solution:**

$$\begin{aligned} \frac{dy}{dx} &= 1 \times e^{-x} + x \times (-1) \times e^{-x} \text{ (using the product rule),} \\ &= e^{-x}(1 - x), \\ &= -e^{-x}(x - 1). \end{aligned}$$

- (ii) Find any stationary points and determine their nature. 2

**Solution:**

$$\frac{dy}{dx} = 0 \text{ when } x = 1.$$

$\therefore$  Maximum at  $(1, 1/e)$ .

	0	1	2
$\frac{dy}{dx}$	1	0	-0.135
	$\nearrow$	$\rightarrow$	$\searrow$

(iii) Prove that  $\frac{d^2y}{dx^2} = e^{-x}(x - 2)$ .

1

**Solution:** 
$$\begin{aligned} \frac{d^2y}{dx^2} &= -(-1)(e^{-x})(x - 1) + 1 \times (-e^{-x}) \text{ (product rule),} \\ &= e^{-x}(x - 1) - e^{-x}, \\ &= e^{-x}(x - 2). \end{aligned}$$

(iv) Show that there is a point of inflexion on this curve and find the co-ordinates of this point.

2

**Solution:**  $\frac{d^2y}{dx^2} = 0$  when  $x = 2$ .  
Change of concavity.


$\therefore$  Inflexion at  $\left(2, \frac{2}{e^2}\right)$ .

	1	2	3
$\frac{d^2y}{dx^2}$	-0.37	0	0.05
	↓		↑

(v) Sketch the curve, showing the co-ordinates of the point of inflexion and any stationary points.

1

**Solution:**



**Question 16 (15 marks)** (use a separate answer booklet)

- (a) The number of DVD copies sold at a store of *The London Olympics 2012 Opening Ceremony* has increased exponentially in accordance with the formula  $N = Ae^{kt}$  where  $t$  is the time in weeks after the Opening Ceremony.

Initially 10 000 copies were sold and the number doubled after two weeks.

- (i) Find the value of  $A$ .

1

**Solution:**

$$\begin{aligned} N &= Ae^{kt}, \\ 10\,000 &= Ae^0, \\ \therefore A &= 10\,000. \end{aligned}$$

- (ii) Calculate the value of  $k$  correct to 3 decimal places.

1

**Solution:**

$$\begin{aligned} 20\,000 &= 10\,000e^{2k}, \\ e^{2k} &= 2, \\ k &= \ln \sqrt{2}, \\ &\approx 0.347. \end{aligned}$$

- (iii) At what rate was the number of copies increasing after four weeks? Answer correct to the nearest whole number.

2

**Solution:**

$$\frac{dN}{dt} = kAe^{kt}.$$

$\therefore$  After 4 weeks,  $\frac{dN}{dt} = 0.347 \times 10\,000e^{4 \times 0.347}$ ,  
 $\approx 13\,863$  (nearest integer).  
 So sales were increasing at 13 863 copies/week.

- (b) Mr B—— borrows  $\$P$  to fund his new Nissan supercar. The term of the loan is 10 years with an interest rate of 6% p.a., monthly reducible. He repays the loan in equal monthly installments of  $\$750$ .

- (i) Show that at the end of  $n$  months, the amount owing is given by  $A = P(1.005)^n - 150\,000(1.005)^n + 150\,000$ .

3

**Solution:** 6% a year is equivalent to a monthly rate of 0.5%.

$$\begin{aligned} \text{Owe after 1 mo.} &= P(1.005) - 750, \\ \text{owe after 2 mo.} &= (P(1.005) - 750)(1.005) - 750, \\ \text{owe after 3 mo.} &= ((P(1.005) - 750)(1.005) - 750)(1.005) - 750, \\ &= P(1.005)^3 - 750(1 + 1.005 + 1.005^2), \\ \text{owe after } n \text{ mo.} &= P(1.005)^n - 750(1 + 1.005 + \dots + 1.005^{n-1}), \\ &= P(1.005)^n - \frac{750(1.005^n - 1)}{1.005 - 1}, \\ &= P(1.005)^n - 150\,000(1.005^n - 1), \\ &= P(1.005)^n - 150\,000(1.005)^n + 150\,000. \end{aligned}$$

- (ii) If at the end of 10 years the loan has been repaid, calculate the amount that he originally borrowed, correct to the nearest dollar. 2

**Solution:**  $0 = P(1.005)^{120} - 150\,000(1.005)^{120} + 150\,000,$   

$$P = \frac{150\,000(1.005^{120} - 1)}{1.005^{120}},$$
  

$$\approx 67\,555.$$
  
 His car cost \$67 555.

- (c) A particle is moving along the  $x$ -axis. The distance of the particle,  $x$  metres from the origin  $O$  is given by the equation  $x = 6t + e^{-4t}$  where  $t$  is time in seconds.

- (i) Write down an expression for velocity of the particle. 1

**Solution:** Let velocity be  $v$ .  

$$x = 6t + e^{-4t},$$
  


$$v = \frac{dx}{dt},$$
  

$$= 6 - 4e^{-4t}.$$

- (ii) Explain why the particle will never come to rest. 2


**Solution:** When  $t = 0$ ,  $v = 2$ ,  
 as  $t \rightarrow \infty$ ,  $v \rightarrow 6$ .  
 Acceleration,  $\frac{dv}{dt} = 16e^{-4t}$  which is always positive.  
 So velocity is always positive and the particle is always accelerating away from the origin and can never come to rest.

- (d)  $A$ ,  $B$ , and  $C$  are the vertices of an isosceles triangle where  $AC = BC$  and right-angled at  $C$ .  $D$  is a point such that  $DB = AB$  and  $\widehat{D\hat{B}A}$  is acute.  $DC \parallel AB$  also. 3



Find, giving reasons, the size of  $\widehat{D\hat{B}C}$ .

**Solution:**



Let  $AC = l = BC$ .

$$l^2 + l^2 = AB^2 \text{ (Pythagoras),}$$

$$\therefore AB = \sqrt{2l^2} = l\sqrt{2},$$

$\widehat{CAB} = \widehat{CBA} = 45^\circ$  (equal base  $\angle$ s of isosc.  $\triangle$ ).

Draw  $BE \perp AB$  and  $DE$  ( $AB \parallel DE$ ),

$$\widehat{CBE} = 45^\circ \text{ (complement of } \widehat{CBA}),$$

$$\widehat{ECB} = 45^\circ \text{ (}\angle \text{ sum } \triangle CEB),$$

$\triangle CEB$  is isosceles (equal base angles),

$$CE = EB \text{ (sides opposite equal } \angle\text{s),}$$

$$2EB^2 = l^2,$$

$$EB = \frac{l}{\sqrt{2}},$$

$$\cos \widehat{EBD} = \frac{l}{\sqrt{2}} \times \frac{1}{l\sqrt{2}},$$

$$= \frac{1}{2},$$

$$\widehat{EBD} = 60^\circ.$$

$$\therefore \widehat{CBD} = \widehat{EBD} - \widehat{CBE},$$

$$= 60^\circ - 45^\circ,$$

$$= 15^\circ.$$

**End of Paper**